

How To Give an MBRT

This document will teach you how to give an MBRT, both oral and rectal, in office.


Prepare the Sample

*NOTE: If any other treatments are being given on the same day, give the MBRT last.

Make sure the blender is thoroughly cleaned before and after use.

Pour room temperature sterile saline into the blender. Then add your donor sample.

*40cc of saline to 2 tablespoons of stool



How To Give an MBRT

Gently Blend the Sample

Attach blender top and blend until you have a smooth mixture. Pulse the blender at first then blend for a short period of time. You do not want to heat up your sample and loose any beneficial bacteria.



How To Give an MBRT

Strain the Sample

Remove blender attachment and attach the grated lid to pour off your mixture, and to keep out any lumps. Pour your mixture into a clean container.



How To Give an MBRT

Draw up your Sample

Draw up the desired amount for your patient. Choose an appropriate catheter and lubricate it before insertion.

- Small dogs and cats = 12-15cc
- Medium dogs = 15-40cc
- Large dog= 40-60cc



How To Give an MBRT

Rectal Ozone

We like to precede the MBRT with Ozone gas (O₃). The tail of the patient must be held down for 10 minutes so they do not expel the gas. Let your patient go outside to poop before the MBRT.




How To Give an MBRT

Inserting the MBRT

Insert your catheter into the patient. Attach syringe with MBRT mixture and inject into the patient. Keep syringe attached to catheter when removing or the MBRT solution will come back out the catheter.

TIP: Leave some air in the syringe to push all of the MBRT through the catheter and into the patient.



How To Give an MBRT

Massage the Abdomen

Massage the abdomen to spread out the MBRT mixture. Try not to let your patient defecate for a few hours, and let the MBRT settle.

